

Mat 1575 HW8 (§3.3)

Simplify the following expressions by writing each one using a single trigonometric function.

126. $4 - 4\sin^2\theta = 4(1 - \sin^2\theta) = 4\cos^2\theta$

128. $a^2 + a^2\tan^2\theta = a^2(1 + \tan^2\theta) = a^2\sec^2\theta$

Integrate using the method of trigonometric substitution. Express the final answer in terms of the variable.

135. $\int \frac{dx}{\sqrt{x^2 - a^2}}$ $x = a\sec\theta \Rightarrow \sec\theta = \frac{x}{a}$
 $dx = a\sec\theta \tan\theta d\theta$
 $= \int \frac{a\sec\theta \tan\theta d\theta}{\sqrt{a^2\sec^2\theta - a^2}} = \int \frac{a\sec\theta \tan\theta d\theta}{\sqrt{a^2\tan^2\theta}} = \int \frac{a\sec\theta \tan\theta d\theta}{|a\tan\theta|}$
 $\Rightarrow x > a \text{ or } x < -a$
 $\Rightarrow a\sec\theta > a \text{ or } a\sec\theta < -a$
 $\Rightarrow \sec\theta > 1 \text{ or } \sec\theta < -1$
 $\Rightarrow 0 \leq \theta < \frac{\pi}{2} \text{ or } \frac{\pi}{2} < \theta \leq \pi$
 $| \tan\theta | = \tan\theta \quad | \tan\theta | = -\tan\theta$


① If $0 \leq \theta < \frac{\pi}{2}$, then $| \tan\theta | = \tan\theta$

(*) $= \int \frac{a\sec\theta \tan\theta}{a\tan\theta} d\theta$


$= \int \sec\theta d\theta$

$= \ln|\sec\theta + \tan\theta| + C$

$= \ln\left|\frac{x}{a} + \frac{\sqrt{x^2 - a^2}}{a}\right| + C$



$\sec\theta = \frac{x}{a}$
 $\tan\theta = \frac{\sqrt{x^2 - a^2}}{a}$



$\sec\theta = \frac{x}{-a}$
 $\tan\theta = \frac{\sqrt{x^2 - a^2}}{-a}$

② If $\frac{\pi}{2} < \theta \leq \pi$, then $| \tan\theta | = -\tan\theta$

(*) $= \int \frac{a\sec\theta \tan\theta}{-a\tan\theta} d\theta$
 $= \int -\sec\theta d\theta$
 $= -\ln|\sec\theta + \tan\theta| + C$
 $= -\ln\left|\frac{x}{a} - \frac{\sqrt{x^2 - a^2}}{a}\right| + C$

$\tan\theta = 3x$
 $x = \frac{1}{3}\tan\theta, dx = \frac{1}{3}\sec^2\theta d\theta$
 $\sqrt{1 + 9x^2} = \sqrt{1 + 9\left(\frac{1}{3}\tan\theta\right)^2}$
 $= \sqrt{1 + \tan^2\theta} = |\sec\theta| = \sec\theta$

137. $\int \frac{dx}{\sqrt{1 + 9x^2}}$
 $= \int \frac{\frac{1}{3}\sec^2\theta}{\sec\theta} d\theta$
 $= \int \frac{1}{3}\sec\theta d\theta = \frac{1}{3}\ln|\sec\theta + \tan\theta| + C$

$\tan\theta = \frac{3x}{1}$



$\sec\theta = \sqrt{1 + 9x^2}$

$1 + 9x^2 \geq 0 \Rightarrow -\infty < x < \infty$
 $\Rightarrow -\infty < \frac{1}{3}\tan\theta < \infty$
 $\Rightarrow -\infty < \tan\theta < \infty$
 $\Rightarrow -\frac{\pi}{2} < \theta < \frac{\pi}{2}$
 $\Rightarrow |\sec\theta| = \sec\theta$

$$= \frac{1}{3} \ln |\sqrt{1-x^2} + 3x| + C$$

139. $\int \frac{dx}{x^2 \sqrt{1-x^2}}$

$x = \sin \theta ; dx = \cos \theta d\theta$
 $\sqrt{1-x^2} = \sqrt{1-\sin^2 \theta} = \sqrt{\cos^2 \theta} = |\cos \theta|$

$= \int \frac{\cos \theta d\theta}{\sin^2 \theta |\cos \theta|} = \int \frac{\cancel{\cos \theta} d\theta}{\sin^2 \theta \cdot \cancel{\cos \theta}} = \int \frac{d\theta}{\sin^2 \theta}$


$= \int \csc^2 \theta d\theta = -\cot \theta + C = -\frac{\sqrt{1-x^2}}{x} + C$

$$1-x^2 \geq 0 \Rightarrow -1 \leq x \leq 1$$

$$-1 \leq \sin \theta \leq 1 \Rightarrow \frac{\pi}{2} \leq \theta \leq \frac{3\pi}{2}$$

Thus, $|\cos \theta| = \cos \theta$

$\sin \theta = \frac{x}{1}$



$\cot \theta = \frac{\sqrt{1-x^2}}{x}$

141. $\int \sqrt{x^2+9} dx$

$x = 3 \tan \theta ; dx = 3 \sec^2 \theta d\theta$
 $\sqrt{x^2+9} = \sqrt{9 \tan^2 \theta + 9} = 3 |\sec \theta|$

$= \int 3 |\sec \theta| d\theta = \int 3 \sec \theta d\theta$


$= 3 \ln |\sec \theta + \tan \theta| + C$

$= 3 \ln \left| \frac{\sqrt{x^2+9}}{3} + \frac{x}{3} \right| + C$

$x^2+9 > 0 \Rightarrow -\infty < x < \infty$
 $\Rightarrow -\infty < 3 \tan \theta < \infty$
 $\Rightarrow -\infty < \tan \theta < \infty$
 $\Rightarrow -\frac{\pi}{2} < \theta < \frac{\pi}{2}$

$\Rightarrow |\sec \theta| = \sec \theta$

$\tan \theta = \frac{x}{3}$



$\sec \theta = \frac{\sqrt{x^2+9}}{3}$


143. $\int \frac{\theta^3 d\theta}{\sqrt{9-\theta^2}}$

$\theta = 3 \sin x ; d\theta = 3 \cos x dx$
 $\sqrt{9-\theta^2} = \sqrt{9-9 \sin^2 x} = 3 |\cos x| = 3 \cos x$

(*) similar statement

$= \int \frac{(3 \sin x)^3}{3 \cos x} \cdot 3 \cos x dx = \int (3 \sin x)^3 dx$

$\sin x = \frac{\theta}{3}$



$\cos x = \frac{\sqrt{9-\theta^2}}{3}$

$= \int 27 \sin^3 x dx = \int 27 (1-\cos^2 x) \sin x dx$

$u = \cos x$
 $du = -\sin x dx$


$= \int -27 (1-u^2) du = \int -27 u^2 + 27 du$

$= -9u^3 + 27u + C = -9 \cos^3 x + 27 \cos x + C$

$= -9 \left(\frac{\sqrt{9-\theta^2}}{3} \right)^3 + 27 \frac{\sqrt{9-\theta^2}}{3} + C$

$= -\frac{(\sqrt{9-\theta^2})^3}{3} + 9\sqrt{9-\theta^2} + C$

147. $\int \frac{dx}{(x^2-9)^{3/2}}$ $x=3\sec\theta, dx=3\sec\theta \cdot \tan\theta d\theta \rightarrow \sec\theta = \frac{x}{3}$
 $(x^2-9)^{3/2} = (\sqrt{x^2-9})^3 = (\sqrt{(3\sec\theta)^2-9})^3$
 $= (3|\tan\theta|)^3$




① If $0 < \theta < \frac{\pi}{2}$, $|\tan\theta| = \tan\theta$, then

$$\int \frac{dx}{(x^2-9)^{3/2}} = \int \frac{3\sec\theta \cdot \tan\theta d\theta}{(3\tan\theta)^3} = \int \frac{\sec\theta}{9\tan^2\theta} d\theta$$

$$= \int \frac{\frac{1}{\cos\theta}}{9 \frac{\sin^2\theta}{\cos\theta}} d\theta = \int \frac{1}{9} \frac{\cos\theta}{\sin^2\theta} d\theta = \int \frac{1}{9} \frac{du}{u^2}$$

$u = \sin\theta$
 $du = \cos\theta d\theta$

$$= -\frac{1}{9} \frac{1}{u} + C = -\frac{1}{9} \frac{1}{\sin\theta} + C = -\frac{1}{9} \frac{x}{\sqrt{x^2-9}} + C$$




② If $\frac{\pi}{2} < \theta \leq \pi$, $|\tan\theta| = -\tan\theta$, then

$$\int \frac{dx}{(x^2-9)^{3/2}} = \int \frac{3\sec\theta \cdot \tan\theta d\theta}{(-3\tan\theta)^3} = \int -\frac{\sec\theta}{9\tan^2\theta} d\theta$$

$$= \dots = \frac{1}{9} \frac{1}{\sin\theta} + C = -\frac{1}{9} \frac{x}{\sqrt{x^2-9}} + C$$

151. $\int \frac{dx}{x^2 \sqrt{x^2+1}}$

$x = \tan\theta, dx = \sec^2\theta d\theta$
 $\sqrt{x^2+1} = \sqrt{\tan^2\theta+1} = |\sec\theta|$
 $= \sec\theta$



$$= \int \frac{\sec^2\theta}{\tan^2\theta \cdot \sec\theta} d\theta$$

$$= \int \frac{\sec\theta}{\tan^2\theta} d\theta = \int \frac{\frac{1}{\cos\theta}}{\frac{\sin^2\theta}{\cos\theta}} d\theta = \int \frac{\cos\theta}{\sin^2\theta} d\theta$$


$u = \sin\theta$
 $du = \cos\theta d\theta$

$$= \int \frac{du}{u^2} = -\frac{1}{u} + C = -\frac{1}{\sin\theta} + C = -\frac{\sqrt{1+x^2}}{x} + C$$

153. $\int (1-x^2)^{3/2} dx$

$x = \sin\theta, dx = \cos\theta d\theta$
 $(1-x^2)^{3/2} = (\sqrt{1-\sin^2\theta})^3 = |\cos\theta|^3 = \dots = \cos^3\theta$
 $\sin\theta = \frac{x}{1}$

$$= \int \cos^3\theta \cdot \cos\theta d\theta = \int \cos^4\theta d\theta = \int \cos^2\theta \cdot \cos^2\theta d\theta$$



$$= \int \left(\frac{1+\cos(2\theta)}{2} \right) \left(\frac{1+\cos(2\theta)}{2} \right) d\theta = \int \frac{1}{4} + \frac{\cos(2\theta)}{2} + \frac{\cos^2(2\theta)}{4} d\theta$$

$$\downarrow$$

$$\cos \theta = \sqrt{1-x^2}$$

$$\theta = \sin^{-1}(x)$$

$$= \int \frac{1}{4} + \frac{\cos(2\theta)}{2} + \frac{1}{4} \cdot \frac{1+\cos(4\theta)}{2} d\theta$$

$$= \int \frac{1}{4} + \frac{\cos(2\theta)}{2} + \frac{1}{8} + \frac{\cos(4\theta)}{8} d\theta = \int \frac{3}{8} + \frac{\cos(2\theta)}{2} + \frac{\cos(4\theta)}{8} d\theta$$

$$= \frac{3}{8}\theta + \frac{\sin(2\theta)}{4} + \frac{\sin(4\theta)}{32} + C$$

$$= \frac{3}{8}\theta + \frac{1}{4} \cdot 2\sin\theta \cdot \cos\theta + \frac{1}{32} 2\sin 2\theta \cos 2\theta + C$$

$$= \frac{3}{8}\theta + \frac{\sin\theta \cos\theta}{2} + \frac{(2\sin\theta \cos\theta)(\cos^2\theta - \sin^2\theta)}{16} + C$$

$$= \frac{3}{8} \sin^{-1}(x) + \frac{1}{2} \cdot x \cdot \sqrt{1-x^2} + \frac{1}{16} (2x\sqrt{1-x^2})(1-x^2-x^2) + C$$

$$= \frac{3}{8} \sin^{-1}(x) + \frac{1}{2} x \cdot \sqrt{1-x^2} + \frac{1}{8} (2x\sqrt{1-x^2})(1-2x^2) + C.$$